

MERCK ANIMAL HEALTH Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 973-937-5557

Website: www.merck-animal-health-usa.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

Prestige[®] V + VEE with Havlogen[®] **Intervet/Merck Animal Health**

ENCEPHALOMYELITIS - RHINOPNEUMONITIS - INFLUENZA VACCINE, EASTERN & WESTERN & VENEZUELAN, KILLED VIRUS, TETANUS TOXOID

A combination of inactivated, concentrated, adjuvanted, tissue culture origin equine encephalomyelitis viruses (Eastern, Western and Venezuelan), equine herpesviruses types 1 and 4 (EHV-1 and EHV-4), equine influenza virus type H3N8 strains (EIV A/Equine 2/ Kentucky/93 American, EIV A/Equine 2/Kentucky/02 American and EIV A/Equine 2/Newmarket/2/93 Eurasian) and tetanus toxoid. Intervet serological data suggest influenza cross protection against certain U.S. and European strains, including Kentucky 91, 93, 99, 02, Ohio 03, Newmarket/1/93, Newmarket/2/93, Suffolk 89 and South Africa 03.

INDICATIONS: For vaccination of healthy horses 6 months of age or older, as an aid in the prevention of disease caused by Eastern, Western and Venezuelan encephalomyelitis viruses and tetanus, as an aid in the control of respiratory disease caused by EIV, EHV-1 and EHV-4 and as an aid in reduction of virus shedding of EIV and EHV-1. Duration of immunity (DOI) has been shown to be at least 6 months for EIV. Foals nursing immune-dams should be vaccinated when maternal antibody levels will allow active immunization.

DOSE: For primary vaccination aseptically administer 1 mL intramuscularly, repeat with a single dose in 3 - 4 weeks. Annual revaccination is recommended. A booster dose can be administered at more frequent intervals based upon individual horse or farm disease risk assessment or any time epidemic conditions exist or are reported. Consult your veterinarian.

CAUTIONS: Store at 2°-7°C (35°-45°F). Shake well before using. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Inject deep into the muscle only. Local reactions may occur at the injection sites in vaccinates. Systemic reactions may also occur in animals following vaccination. If allergic reaction occurs, treat with epinephrine. Care should be used to avoid injecting your own fingers, hands, or other body parts. Accidental injection can cause serious local reactions. Contact a physician immediately if an accidental injection occurs.

ANTIBIOTIC: Gentamicin.

PRESERVATIVE: Thimerosal.

Adjuvant: Intervet's Proprietary Technology

FOR ANIMAL USE ONLY

Intervet Inc., Merck Animal Health, division of Intervet Inc., Omaha, NE 68103, USA

U.S. Vet Lic. No 165A

1 800 521-5767 (USA)

1 866 683-7838 (CAN)

For patent information:

<http://www.merck.com/product/patent/home.html>

		Code	
10 Doses	10 x 1 mL Syringes with Separate Sterile Needles	030313	145893-02
10 Doses	10 mL	032887	139888-03

CPN: 1047414.3

