

IRON DEXTRAN-100

Indications:

Injectable Iron 100 mg is a sterile, injectable solution for use in the prevention or treatment of baby pig anemia due to iron deficiency.

Approved For Use On:

Swine

Active Ingredients:

Elemental iron....100mg/mL

Product Advantages:

- Reduces losses due to iron deficiency.
- Helps achieve optimum levels of iron necessary for good growth and overall good health.
- Safe.
- Quick absorption.
- Good syringeability.

Dosage and Administration:

For the prevention of anemia due to iron deficiency, administer an intramuscular injection of 100mg of elemental iron (1mL) at 2 to 4 days of age.

For the treatment of anemia due to iron deficiency, administer an intramuscular injection of 100mg of elemental iron (1mL) at the first signs of anemia. May be repeated in approximately 10 days.


Packaging:

100mL bottle, 12/case UPC # 7-45801-11030-4



IRON DEXTRAN-100

DIRECTIONS FOR USE: Disinfect rubber stopper of vial and injection site with a cotton ball moistened with rubbing alcohol or other suitable disinfectant. Use a small needle (20 gauge, 5/8 inch) and syringe that have been sterilized by boiling in water for 20 minutes. Injection should be made intramuscularly into the back of the ham (see diagram below). Apply tension to the skin at the injection site by pulling skin and fat downward with pressure from the thumb or fingers. Inject into the muscle to a depth of approximately 1/2 inch. Use a clean, sterile needle and syringe for each pig to avoid spreading infection.



PRECAUTIONS: If disease or abnormal condition is suspected, consult veterinarian before administration.

SIDE EFFECTS: Occasionally pigs may exhibit a reaction to iron injection, clinically characterized by prostration and muscular weakness. In extreme cases, death may result.

NOTICE: The intramuscular injection of organic iron preparations into pigs beyond 4 weeks of age may cause staining of muscle tissue.

HOW SUPPLIED: 100 mL multiple dose vials.

Rev. 12-08

I-0900-04