

MERCK ANIMAL HEALTH

Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

Customer Service: 800-521-5767
Order Desk: 800-648-2118
Technical Service (Companion Animal): 800-224-5318
Technical Service (Livestock): 800-211-3573
Fax: 973-937-5557
Website: www.merck-animal-health-usa.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

SAFE-GUARD® DEWORMER FOR GOATS

Intervet/Merck Animal Health

(fenbendazole)

Suspension 10% (100 mg/mL)

NOT FOR HUMAN USE

RESIDUE WARNINGS: Goats must not be slaughtered for food within 6 days following treatment. Because a withdrawal time in milk has not been established, do not use in lactating goats.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Keep this and all medication out of the reach of children.

INDICATIONS:

Goats - 2.3 mg/lb (5 mg/kg) body weight for the removal and control of: Stomach worms (adults): *Haemonchus contortus* and *Teladorsagia circumcincta*.

DIRECTIONS:

Determine the proper dose according to estimated body weight. Administer orally. The recommended dose of 2.3 mg/lb (5 mg/kg) is achieved when 2.3 mL of the drug are given for each 100 lb body weight.

EXAMPLES: Goats:

Body Weight	Amount
25 lb	0.6 mL
50 lb	1.2 mL
75 lb	1.7 mL
100 lb	2.3 mL
125 lb	2.9 mL

Under conditions of continued exposure to parasites, retreatment may be needed after 4 to 6 weeks.

Manufactured by: DPT Laboratories, San Antonio, TX 78215

Distributed by: **Intervet Inc.**, Millsboro, DE 19966

www.intervetusa.com

Store at or below 25°C (77°F). Protect from freezing. Shake well before use. Restricted drug (California) - Use only as directed.

NADA # 128-620, Approved by FDA

125 mL (4.2 fl oz)	697600-B, 697650-B
--------------------	--------------------

CPN: 1047469.1