

MERCK ANIMAL HEALTH

Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 973-937-5557

Website: www.merck-animal-health-usa.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

safe-guard[®] Canine Dewormer Intervet/Merck Animal Health

(fenbendazole)

For dogs only, 6 weeks and older.

Drug Facts:

Active ingredients (in each dosage unit): Fenbendazole Granules 22.2% (222 mg/g)

Uses: For the treatment and control of:

- Tapeworms (*Taenia pisiformis*)
- Roundworms (*Toxocara canis*, *Toxascaris leonina*)
- Hookworms (*Ancylostoma caninum*, *Uncinaria stenocephala*)
- Whipworms (*Trichuris vulpis*)

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

HUMAN WARNING: KEEP OUT OF REACH OF CHILDREN.

Directions: The daily dose for Safe-Guard[®] Canine is 50 mg/kg (22.7 mg/lb) of body weight. You should weigh your dog to make sure you are using the right dose; i.e. correct gram amount and appropriate number of pouches.

Your dog must be treated with the following dose for 3 days in a row.

Dog Weight (lbs)	Pouch Size (Daily Dose)
10	1 gram
11 to 20	2 gram
21 to 30*	1 gram + 2 gram
31 to 40	4 gram
41 to 50*	1 gram + 4 gram
51 to 60*	2 gram + 4 gram
61 to 80*	Two 4 gram
Over 80*	* Combine pouches to obtain recommended daily dosage.

Mix daily dose with a small amount of the usual food. Make sure your dog eats all the medicated food. You may need to moisten dry dog food to aid mixing.

Repeat the dose for 3 days in a row.

Recommended Deworming Schedule: Schedule may vary depending on the climate where you live and the activity of your dog. The following schedule should be used as a general guideline.

- Weaned pups (6-8 weeks of age) should be dewormed at 6, 8, 10 and 12 weeks of age.
- Treat the dam at the same time as the puppies.
- Dogs over 6 months of age should be dewormed at least twice each year.

Diagnosis of Parasites: Consult your veterinarian for help in the diagnosis, treatment and control of parasitism. Specific diagnoses require laboratory

testing. Dogs can become infected with several different kinds of tapeworms. Safe-Guard® Canine only kills *Taenia* species of tapeworms. If you continue to see tapeworm segments in your dog's stool after treatment with Safe-Guard® Canine, consult your veterinarian.

Roundworms and hookworms may pose serious human health risks; regular deworming may help reduce those risks.

Product Use: Safe-Guard® Canine is safe for use in puppies 6 weeks or older and adult dogs, including pregnant bitches. Do not deworm a dog or puppy that is sick. Consult your veterinarian for diagnosis of the illness.

Adverse Reactions: In U.S. clinical studies 3 of 240 dogs (about 1% of the treated dogs) had vomiting associated with the use of the product.

Storage: Store at controlled room temperature 68-77°F (20-25°C).

Questions? Comments?

To obtain product information including material safety data sheets (MSDS) or to report a suspected adverse reaction, call **1-800-441-8272**.

Distributed by: Intervet Inc (d/b/a Merck Animal Health), Summit, NJ 07901

NADA #121-473, Approved by FDA

Made in Austria

www.Safe-Guard.com

Net Weight:		
3 grams	Three 1 gram pouches for a 3-day treatment. Each pouch treats 10 lbs.	114853 R1 446100-D
6 grams	Three 2 gram pouches for a 3-day treatment. Each pouch treats 20 lbs.	110895 R1 446200-D
12 grams	Three 4 gram pouches for a 3-day treatment. Each pouch treats 40 lbs.	119438 R1 446400-D

CPN: 1047429.2